

Simplified HCV Treatment for Treatment-Naïve Adults Without Cirrhosis Screening Questionnaire

Date: _____

Patient Name: _____ ID#: _____ DOB: _____

Provider (or designee) please complete Sections 1 and 2 and return to The Hepatitis C Committee

Section 1 – Review for Contraindications

- Has confirmed diagnosis of Hepatitis C
- Has FibroScan Score of < 12.5 kPa
- Has not previously been treated for Hep C
- No prior hypersensitivity to Hep C treatment
- Not on Hospice or have a life expectancy of < 1 year. *Illnesses that could be remediated by treatment of Hep C does not disqualify patient from receiving the therapy.
- Patient will have sufficient time to complete treatment (Typically 12 weeks. Confirm with patient’s counselor of release plan.)
- Patient is willing to be treated for Hep C with a once daily oral medication for 8 to 12 weeks

Comments:

Section 2 – Review Documentation Completion

- IDPH Sexual History completed
- Review and signed informed consent (**HSF 912 F-2**)

Section 3 – Provider – Required Exam and Testing

- Physician Encounter to rule out clinical signs or symptoms of cirrhosis and educate patient on proper administration of medication, adherence, and prevention of reinfection.
- Laboratory Testing Within the Last 6 Months:
 - CBC
 - CMP (LFT’s and BMP)
- Laboratory Testing Any Time Prior to Therapy:
 - Quantitative HCV RNA (HCV viral load)
 - HIV antigen/antibody test
 - Hepatitis B surface antigen
 - HCV Genotype
- Before Initiating Therapy:
 - Serum pregnancy testing and counseling about pregnancy risks of HCV medication should be offered to women of child bearing age.
 - A negative pregnancy test should be confirmed prior to initiating therapy.

Section 4 – Provider Educate and Consent

- Review treatment recommendations with patient.
- Enter diagnosis “Hepatitis Treatment” in Medical ICON

Distribution: Original HSF-912 F-1, F-2 and IDPH history scanned into Medical ICON, Copies to Hepatitis C Committee